

兔抗 ACSBG2 多克隆抗体

中文名称：兔抗 ACSBG2 多克隆抗体

英文名称：Anti-ACSBG2 rabbit polyclonal antibody

别名：BGR, BRGL, PRDNY3, PRD-NY3

相关类别：一抗

储存：冷冻（-20℃）

抗原：ACSBG2

宿主：Rabbit

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

| | |
|------------------------------------|--|
| Background: | Long-chain-fatty-acid—CoA ligase ACSBG2 is an enzyme that in humans is encoded by the ACSBG2 gene. Mediates activation of long-chain fatty acids for both synthesis of cellular lipids, and degradation via beta-oxidation. Able to activate long-chain fatty acids. Also able to activate very long-chain fatty acids, however, the relevance of such activity is unclear in vivo. Has increased ability to activate oleic and linoleic acid. May play a role in spermatogenesis. |
| Applications: | ELISA, IHC |
| Name of antibody: | ACSBG2 |
| Immunogen: | Fusion protein of human ACSBG2 |
| Full name: | acyl-CoA synthetase bubblegum family member 2 |
| Synonyms : | BGR, BRGL, PRTDNY3, PRTD-NY3 |
| SwissProt: | Q5FVE4 |
| ELISA Recommended dilution: | 500-5000 |
| IHC positive control: | Human stomach cancer |
| IHC Recommend dilution: | 50-100 |

