

兔抗 FZR1 多克隆抗体

- 中文名称：兔抗 FZR1 多克隆抗体
- 英文名称：Anti-FZR1 rabbit polyclonal antibody
- 别名：FZR; CDH1; FZR2; HCDH; HCDH1; CDC20C
- 相关类别：一抗
- 抗原：FZR1
- 储存：冷冻（-20℃）避光
- 宿主：Rabbit
- 反应种属：Human, Mouse, Rat
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

技术规格

Background:	Substrate-specific adapter for the anaphase promoting complex/cyclosome (APC/C) E3 ubiquitin-protein ligase complex. Associates with the APC/C in late mitosis, in replacement of CDC20, and activates the APC/C during anaphase and telophase. The APC/C remains active in degrading substrates to ensure that positive regulators of the cell cycle do not accumulate prematurely. At the G1/S transition FZR1 is phosphorylated, leading to its dissociation from the APC/C. Following DNA damage, it is required for the G2 DNA damage checkpoint: its dephosphorylation and reassociation with the APC/C leads to the ubiquitination of PLK1, preventing entry into mitosis. Acts as an adapter for APC/C to target the DNA-end resection factor RBBP8/CtIP for ubiquitination and subsequent proteasomal degradation. Through the regulatio
--------------------	---

	n of RBBP8/CtIP protein turnover, may play a role in DNA damage response, favoring DNA double-strand repair through error-prone non-homologous end joining (NHEJ) over error-free, RBBP8-mediated homologous recombination (HR) (PubMed:25349192).
Applications:	WB, IHC
Name of antibody:	FZR1
Immunogen:	Fusion protein of human FZR1
Full name:	fizzy and cell division cycle 20 related 1
Synonyms:	FZR; CDH1; FZR2; HCDH; HCDH1; CDC20C
SwissProt:	Q9UM11
IHC positive control:	Human breast cancer and Human esophagus cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	55 kDa
WB Positive control:	Human placenta tissue
WB Recommended dilution:	500-2000



