

兔抗 AGA 多克隆抗体

中文名称：兔抗 AGA 多克隆抗体

英文名称：Anti-AGA rabbit polyclonal antibody

别名：GA; AGU; ASRG

相关类别：一抗

抗原：AGA

储存：冷冻（-20℃）

宿主：Rabbit

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

| | |
|------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Background: | Aspartylglucosaminidase is involved in the catabolism of N-linked oligosaccharides of glycoproteins. It cleaves asparagine from N-acetylglucosamines as one of the final steps in the lysosomal breakdown of glycoproteins. The lysosomal storage disease aspartylglycosaminuria is caused by a deficiency in the AGA enzyme. Alternatively spliced transcript variants have been identified. |
| Applications: | ELISA, WB, IHC |
| Name of antibody: | AGA |
| Immunogen: | Fusion protein of human AGA |
| Full name: | Aspartylglucosaminidase |
| Synonyms : | GA; AGU; ASRG |
| SwissProt: | P20933 |
| ELISA Recommended dilution: | 1000-2000 |
| IHC positive control: | Human breast cancer and Human lung cancer |
| IHC Recommend dilution: | 15-50 |
| WB Predicted band size: | 37 kDa |
| WB Positive control: | Mouse kidney tissue, human testis and thyroid cancer tissue |
| WB Recommended dilution: | 200-1000 |



