

兔抗 AKT1 多克隆抗体

中文名称: 兔抗 AKT1 多克隆抗体

英文名称: Anti-AKT1 rabbit polyclonal antibody

别 名: AKT, PKB, RAC, PRKBA, PKB-ALPHA, RAC-ALPHA

相关类别: 一抗

抗 原: AKT1

储 存: 冷冻(-20℃)

宿 主: Rabbit

反应种属: Human, Mouse, Rat

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

KT2 are activated by platelet-derived growth factor. The activation is rapid and specific, and it is abrogated by mutations in the pleckstrin homology domain of AKT1.

Background:

It was shown that the activation occurs through phosph atidylinositol 3-kinase. In the developing nervous syste m AKT is a critical mediator of growth factor-induced n euronal survival. Survival factors can suppress apoptosis in a transcription-independent manner by activating the serine/threonine kinase AKT1, which then phosphorylate

The serine-threonine protein kinase encoded by the AK T1 gene is catalytically inactive in serum-starved primar y and immortalized fibroblasts. AKT1 and the related A

s and inactivates components of the apoptotic machine



全国订货电话 4008-723-722

| | ry. Mutations in this gene have been associated with the Proteus syndrome. Multiple alternatively spliced transcript variants have been found for this gene. |
|-----------------------------|--|
| Applications: | ELISA, IHC |
| Name of antibody: | AKT1 |
| Immunogen: | Fusion protein of human AKT1 |
| Full name: | V-akt murine thymoma viral oncogene homolog 1 |
| Synonyms: | AKT, PKB, RAC, PRKBA, PKB-ALPHA, RAC-ALPHA |
| SwissProt: | P31749 |
| ELISA Recommended dilution: | 1000-5000 |
| IHC positive control: | Human colon cancer and human tonsil |
| IHC Recommend dilution: | 25-100 |

