

## 兔抗 ARID2 多克隆抗体

中文名称：兔抗 ARID2 多克隆抗体

英文名称：Anti-ARID2 rabbit polyclonal antibody

别名：p200; BAF200

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：ARID2

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

|                          |   |
|--------------------------|---|
| <b>Background:</b>       | ARID2 is a subunit of the PBAF chromatin-remodeling complex (see BAF180; MIM 606083), which facilitates ligand-dependent transcriptional activation by nuclear receptors. Involved in transcriptional activation and repression of select genes by chromatin remodeling (alteration of DNA-nucleosome topology). Required for the stability of the SWI/SNF chromatin remodeling complex SWI/SNF-B (PBAF). |
| <b>Applications:</b>     | ELISA, WB, IHC  |
| <b>Name of antibody:</b> | ARID2   |
| <b>Immunogen:</b>        | Synthetic peptide of human ARID2  |
| <b>Full name:</b>        | AT rich interactive domain 2 (ARID, RFX-like)   |
| <b>Synonyms:</b>         | p200; BAF200  |
| <b>SwissProt:</b>        | Q68CP9  |

|                                    |   |
|------------------------------------|---|
| <b>ELISA Recommended dilution:</b> | 5000-10000                                    |
| <b>IHC positive control:</b>       | Human thyroid cancer and human ovarian cancer |
| <b>IHC Recommend dilution:</b>     | 25-100  |
| <b>WB Predicted band size:</b>     | 197 kDa                                       |
| <b>WB Positive control:</b>        | TM4 cell                                      |
| <b>WB Recommended dilution:</b>    | 200-1000                                      |



