

兔抗 BTN2A2 多克隆抗体

中文名称：兔抗 BTN2A2 多克隆抗体

英文名称： Anti-BTN2A2 rabbit polyclonal antibody

别名： BTF2; BT2.2

相关类别： 一抗

抗原： BTN2A2

储存： 冷冻（-20℃）

宿主： Rabbit

反应种属： Human

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

| | |
|--------------------------|---|
| Background: | Butyrophilin is the major protein associated with fat droplets in the milk. This gene is a member of the BTN2 subfamily of genes, which encode proteins belonging to the butyrophilin protein family. The gene is located in a cluster on chromosome 6, consisting of seven genes belonging to the expanding B7/butyrophilin-like group, a subset of the immunoglobulin gene superfamily. The encoded protein is a type I receptor glycoprotein involved in lipid, fatty-acid and sterol metabolism. Several alternatively spliced transcript variants encoding different isoforms have been found for this gene. |
| Applications: | ELISA, IHC |
| Name of antibody: | BTN2A2 |

| | |
|------------------------------------|--------------------------------------|
| Immunogen: | Fusion protein of human BTN2A2 |
| Full name: | butyrophilin, subfamily 2, member A2 |
| Synonyms : | BTF2; BT2.2 |
| SwissProt: | Q8WVV5 |
| ELISA Recommended dilution: | 2000-5000 |
| IHC positive control: | Human thyroid cancer |
| IHC Recommend dilution: | 25-100 |

