

## 兔抗 CCL2 多克隆抗体

中文名称: 兔抗 CCL2 多克隆抗体

英文名称: Anti-CCL2 rabbit polyclonal antibody

别名: HC11, MCAF, MCP1, MCP-1, SCYA2, GDCF-2, SMC-CF, HSMCR30

相关类别: 一抗

储 存: 冷冻(-20℃)

抗原: CCL2

宿 主: Rabbit

反应种属: Human

标记物: Unconjugate

克隆类型: rabbit polyclonal

## 技术规格

nd inflammatory processes. The protein encoded by th is gene is structurally related to the CXC subfamily of cytokines. Members of this subfamily are characterized by two cysteines separated by a single amino acid. Thi s cytokine displays chemotactic activity for monocytes and basophils but not for neutrophils or eosinophils. It has been implicated in the pathogenesis of diseases c haracterized by monocytic infiltrates, like psoriasis, rhe

okine receptors CCR2 and CCR4.

This gene is one of several cytokine genes clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory a

umatoid arthritis and atherosclerosis. It binds to chem



## 全国订货电话 4008-723-722

Applications:	ELISA, IHC
Name of antibody:	CCL2
Immunogen:	Synthetic peptide of human CCL2
Full name:	C-C motif chemokine 2
Synonyms :	HC11, MCAF, MCP1, MCP-1, SCYA2, GDCF-2, SMC-CF, HSMCR30
SwissProt:	P13500
ELISA Recommended dilution:	500-5000
IHC positive control:	Human breast cancer
IHC Recommend dilution:	25-50

