

兔抗 CCL18 多克隆抗体

中文名称：兔抗 CCL18 多克隆抗体

英文名称：Anti-CCL18 rabbit polyclonal antibody

别名：CKb7; PARC; AMAC1; DCCK1; MIP-4; AMAC-1; DC-CK1; SCYA18

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：CCL18

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

This antimicrobial gene is one of several Cys-Cys (C C) cytokine genes clustered on the q arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for naive T cells, CD4+ and CD8+ T cells and nonactivated lymphocytes, but not for monocytes or granulocytes. This chemokine attracts naive T lymphocytes toward dendritic cells and activated macrophages in lymph nodes. It may play a role in both humoral and cell-mediated immunity responses.

Applications:	ELISA, WB
Name of antibody:	CCL18
Immunogen:	Synthetic peptide of human CCL18
Full name:	chemokine (C-C motif) ligand 18 (pulmonary and activation-regulated)
Synonyms:	CKb7; PARC; AMAC1; DCCK1; MIP-4; AMAC-1; DC-C K1; SCYA18
SwissProt:	P55774
ELISA Recommended dilution:	5000-10000
WB Predicted band size:	10 kDa
WB Positive control:	Human lung tissue lysate
WB Recommended dilution:	500-2000

