

兔抗 CD70 多克隆抗体

中文名称：兔抗 CD70 多克隆抗体

英文名称：Anti-CD70 rabbit polyclonal antibody

别名：CD70; CD27L; CD27LG; TNFSF7; CD70 antigen; Tumor necrosis factor ligand superfamily member 7

相关类别：一抗

储存：冷冻（-20℃）避光

宿主：Rabbit

抗原：CD70

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

| | |
|--------------------------|--|
| Background: | The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T and B lymphocytes. It induces proliferation of costimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin synthesis. |
| Applications: | WB |
| Name of antibody: | CD70 |
| Immunogen: | Fusion protein of human CD70 |

| | |
|---------------------------------|--|
| Full name: | CD70 molecule |
| Synonyms : | CD70; CD27L; CD27LG; TNFSF7; CD70 antigen; Tumor necrosis factor ligand superfamily member 7 |
| SwissProt: | P32970 |
| WB Predicted band size: | 21 kDa |
| WB Positive control: | NCI-H460 cells |
| WB Recommended dilution: | 500-2000 |



