

兔抗 CDA 多克隆抗体

中文名称：兔抗 CDA 多克隆抗体

英文名称：Anti-CDA rabbit polyclonal antibody

别名：cytidine deaminase; CDD

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：CDA

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

| | |
|--------------------------|--|
| Background: | This gene encodes an enzyme involved in pyrimidine salvaging. The encoded protein forms a homotetramer that catalyzes the irreversible hydrolytic deamination of cytidine and deoxycytidine to uridine and deoxyuridine, respectively. It is one of several deaminases responsible for maintaining the cellular pyrimidine pool. Mutations in this gene are associated with decreased sensitivity to the cytosine nucleoside analogue cytosine arabinoside used in the treatment of certain childhood leukemias. |
| Applications: | ELISA, WB |
| Name of antibody: | CDA |
| Immunogen: | Full length fusion protein |
| Full name: | cytidine deaminase |

| | |
|------------------------------------|-----------------------------|
| Synonyms: | CDD |
| SwissProt: | P32320 |
| ELISA Recommended dilution: | 5000-10000 |
| WB Predicted band size: | 16 kDa |
| WB Positive control: | Human stomach tissue lysate |
| WB Recommended dilution: | 200-1000 |

