

兔抗 WNT10A 多克隆抗体

- 中文名称: 兔抗 WNT10A 多克隆抗体
- 英文名称: Anti-WNT10A rabbit polyclonal antibody
- 别 名: OODD; SSPS; STHAG4
- 相关类别: 一抗
- 储 存: 冷冻(-20℃)
- 宿 主: Rabbit
- 抗 原: WNT10A
- 反应种属: Human, Mouse
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

d T s re 0 fa fa d g c c T	The WNT gene family consists of structurally relate d genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesi and in several developmental processes, including egulation of cell fate and patterning during embry ogenesis. This gene is a member of the WNT gene amily. It is strongly expressed in the cell lines of p omyelocytic leukemia and Burkitt's lymphoma. In a ddition, it and another family member, the WNT6 gene, are strongly coexpressed in colorectal cancer cell lines. The gene overexpression may play key ro es in carcinogenesis through activation of the WN T-beta-catenin-TCF signaling pathway. This gene an d the WNT6 gene are clustered in the chromosom
-------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

reagent.com

全国订货电话 4008-723-722

	e 2q35 region.
Applications:	ELISA, IHC
Name of antibody:	WNT10A
Immunogen:	Synthetic peptide of humanWNT10A
Full name:	wingless-type MMTV integration site family, membe r 10A
Synonyms :	OODD; SSPS; STHAG4
SwissProt:	Q9GZT5
ELISA Recommended dilution:	1000-2000
IHC positive control:	Human liver cancer
IHC Recommend dilution:	10-50

