

## 兔抗 HSD17B4 多克隆抗体

中文名称: 兔抗 HSD17B4 多克隆抗体

英文名称: Anti-HSD17B4 rabbit polyclonal antibody

别名: DBP; MFE-2; MPF-2; PRLTS1; SDR8C1

相关类别: 一抗

储存: 冷冻 (-20℃)

抗原: HSD17B4

宿主: Rabbit

反应种属: Human, Mouse, Rat

标记物: Unconjugate

克隆类型: rabbit polyclonal

### 技术规格

<b>Background:</b>	The protein encoded by this gene is a bifunctional enzyme that is involved in the peroxisomal beta-oxidation pathway for fatty acids. It also acts as a catalyst for the formation of 3-ketoacyl-CoA intermediates from both straight-chain and 2-methyl-branched-chain fatty acids. Defects in this gene that affect the peroxisomal fatty acid beta-oxidation activity are a cause of D-bifunctional protein deficiency (DBPD). An apparent pseudogene of this gene is present on chromosome 8.
<b>Applications:</b>	ELISA, WB, IHC
<b>Name of antibody:</b>	HSD17B4
<b>Immunogen:</b>	Fusion protein of human HSD17B4
<b>Full name:</b>	hydroxysteroid (17-beta) dehydrogenase 4

<b>Synonyms :</b>	DBP; MFE-2; MPF-2; PRLTS1; SDR8C1
<b>SwissProt:</b>	P51659
<b>ELISA Recommended dilution:</b>	2000-5000
<b>IHC positive control:</b>	Human colon cancer and human cervical cancer
<b>IHC Recommend dilution:</b>	50-200
<b>WB Predicted band size:</b>	80 kDa
<b>WB Positive control:</b>	Huvec and K562 cells
<b>WB Recommended dilution:</b>	500-2000



