

兔抗 IFIH1 多克隆抗体

中文名称: 兔抗 IFIH1 多克隆抗体

英文名称: Anti-IFIH1 rabbit polyclonal antibody

别名: AGS7; Hlcd; MDA5; MDA-5; RLR-2; IDDM19

抗原: IFIH1

储 存: 冷冻(-20℃) 避光

宿 主: Rabbit

反应种属: Human

相关类别: 一抗

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

a number of cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. This gene encodes a DEAD box protein that is upregulated in response to treatment with beta-interferon and a protein kinase C-activating compound, mezerein. Irreversible reprogramming of melanomas can be achieved by treatment with both these ag

ents; treatment with either agent alone only achieves reversible diff erentiation. Genetic variation in this gene is associated with diabet

DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in

Background:



全国订货电话 4008-723-722

	es mellitus insulin-dependent type 19.
Applications:	WB
Name of antibody:	IFIH1
Immunogen:	Synthesized peptide derived from C-terminal of human IFIH1.
Full name:	interferon induced with helicase C domain 1
Synonyms:	AGS7; Hlcd; MDA5; MDA-5; RLR-2; IDDM19
SwissProt:	Q9BYX4
WB Predicted band s ize:	117 kDa
WB Positive control:	Jurkat cells lysate
WB Recommended d ilution:	500-3000

