

## 兔抗 KCNK17 多克隆抗体

中文名称: 兔抗 KCNK17 多克隆抗体

英文名称: Anti-KCNK17 rabbit polyclonal antibody

别名: TALK2; TASK4; TALK-2; TASK-4; K2p17.1

相关类别: 一抗

储存: 冷冻 (-20℃)

宿主: Rabbit

抗原: KCNK17

反应种属: Human

标记物: Unconjugate

克隆类型: rabbit polyclonal

### 技术规格

<b>WB Recommended dilution:</b>	200-1000
<b>WB Positive control:</b>	LO2 cells
<b>WB Predicted band size:</b>	37 kDa
<b>IHC Recommend dilution:</b>	100-300
<b>IHC positive control:</b>	Human colon cancer and Human thyroid cancer
<b>ELISA Recommended dilution:</b>	1000-2000
<b>SwissProt:</b>	Q96T54
<b>Synonyms :</b>	TALK2; TASK4; TALK-2; TASK-4; K2p17.1
<b>Full name:</b>	potassium channel, subfamily K, member 17

<b>Immunogen:</b>	Fusion protein of human KCNK17
<b>Name of antibody:</b>	KCNK17
<b>Applications:</b>	ELISA, WB, IHC
<b>Background:</b>	The protein encoded by this gene belongs to the family of potassium channel proteins containing two pore-forming P domains. This channel is an open rectifier which primarily passes outward current under physiological K <sup>+</sup> concentrations. This gene is activated at alkaline pH. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.



