

小鼠抗 PSMA6 单克隆抗体

中文名称: 小鼠抗 PSMA6 单克隆抗体

英文名称: Anti-PSMA6 mouse monoclonal antibody

别名: IOTA; p27K; PROS27

抗原: PSMA6

储存: 冷冻 (-20℃) 避光

宿主: Mouse

反应种属: Human , Dog , Rat , Monkey , Mouse

相关类别: 一抗

标记物: Unconjugate

克隆类型: mouse monoclonal

技术规格

Background:	The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. A pseudogene has been identified on the Y chromosome. [provided by RefSeq].
Applications:	WB, IHC, FC

Name of antibody:	PSMA6
Immunogen:	Fusion protein of human PSMA6
Full name:	proteasome (prosome, macropain) subunit, alpha type, 6 (PSMA6), transcript variant 1
Synonyms:	IOTA; p27K; PROS27
SwissProt:	P60900
IHC positive control:	carcinoma of human lung tissue and adenocarcinoma of human ovary tissue; carcinoma of human bladder tissue and carcinoma of human prostate tissue
IHC Recommend dilution:	30-150
WB Predicted band size:	27 kDa
WB Positive control:	HepG2, Hela, SVT2, A549, COS7, Jurkat, MDCK, PC12, MCF-7 cell lysates
WB Recommended dilution:	500-2000