

小鼠抗 SDCBP 单克隆抗体

中文名称: 小鼠抗 SDCBP 单克隆抗体

英文名称: Anti-SDCBP mouse monoclonal antibody

别 名: MDA-9; MDA9; ST1; SYCL; TACIP18

储 存: 冷冻 (-20℃) 避光

抗 原: SDCBP

宿 主: Mouse

反应种属: Human

相关类别: 一抗

标 记 物: Unconjugate

克隆类型: mouse monoclonal

技术规格

Background:	The protein encoded by this gene was initially identified as a molecule linking syndecan-mediated signaling to the cytoskele ton. The syntenin protein contains tandemly repeated PDZ do mains that bind the cytoplasmic, C-terminal domains of a vari ety of transmembrane proteins. This protein may also affect c ytoskeletal-membrane organization, cell adhesion, protein trafficking, and the activation of transcription factors. The protein is primarily localized to membrane-associated adherens junctions and focal adhesions but is also found at the endoplasmic r eticulum and nucleus. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by Ref Seq].
Applications:	WB, IHC



全国订货电话 4008-723-722

_	
Name of antibody:	SDCBP
Immunogen:	Fusion protein of human SDCBP
Full name:	syndecan binding protein (syntenin) (SDCBP), transcript variant
Synonyms:	MDA-9; MDA9; ST1; SYCL; TACIP18
SwissProt:	O00560
IHC positive control:	adenocarcinoma of human ovary tissue and adenocarcinoma of human endometrium tissue; human liver tissue
IHC Recommend dilution:	30-150
WB Predicted band size:	32 kDa
WB Positive control:	HepG2, Hela, A549, Jurkat, MCF-7 cell lysates
WB Recommended dilutio n:	500-2000

