

小鼠抗 TANK 单克隆抗体

中文名称:	小鼠抗 TANK 单克隆抗体
英文名称:	Anti-TANK mouse monoclonal antibody
别名:	TRAF family member associated NFKB activator; ITRAF; TRAF2; I-TRAF
相关类别:	一抗
储存:	冷冻 (-20℃)
宿主:	Mouse
抗原:	TANK
反应种属:	Human, Rat
标记物:	Unconjugate
克隆类型:	mouse monoclonal

技术规格

Background:	The TRAF (tumor necrosis factor receptor-associated factor) family of proteins associate with and transduce signals from members of the tumor necrosis factor receptor superfamily. The protein encoded by this gene is found in the cytoplasm and can bind to TRAF1, TRAF2, or TRAF3, thereby inhibiting TRAF function by sequestering the TRAFs in a latent state in the cytoplasm. For example, the protein encoded by this gene can block TRAF2 binding to LMP1, the Epstein-Barr virus transforming protein, and inhibit LMP1-mediated NF-kappa-B activation. Three alternatively spliced transcript variants encoding different isoforms have been found for this gene.
Applications:	WB, IHC

Name of antibody:	TANK
Immunogen:	Fusion protein of human TANK
Full name:	TRAF family member associated NFKB activator
Synonyms:	ITRAF; TRAF2; I-TRAF
SwissProt:	Q92844
IHC positive control:	Human colorectal cancer and Human appendix tissue; Human colon tissue
IHC Recommend dilution:	200-500
WB Predicted band size:	48 KD
WB Positive control:	HepG2, Hela, A549, MCF7 cell lysates
WB Recommended dilution:	500-2000