

## 小鼠抗 TBCC 单克隆抗体

中文名称: 小鼠抗 TBCC 单克隆抗体

英文名称: Anti-TBCC mouse monoclonal antibody

别 名: CFC

储 存 冷冻(-20℃) 避光

抗 原 TBCC

宿 主 Mouse

反应种属 Human , Rat , Monkey , Mouse

相关类别 一抗

标记物 Unconjugate

克隆类型 mouse monoclonal

## 技术规格

Background:	Cofactor C is one of four proteins (cofactors A, D, E, and C) involved in the pathway leading to correctly folded beta-tub ulin from folding intermediates. Cofactors A and D are belie ved to play a role in capturing and stabilizing beta-tubulin i ntermediates in a quasi-native confirmation. Cofactor E bind s to the cofactor D/beta-tubulin complex; interaction with c ofactor C then causes the release of beta-tubulin polypeptid es that are committed to the native state. [provided by Ref Seq].
Applications:	WB, IHC, IF
Name of antibody:	TBCC
Immunogen:	Fusion protein of human TBCC
Full name:	tubulin folding cofactor C (TBCC)



## 全国订货电话 4008-723-722

Synonyms:	CFC
SwissProt:	Q15814
IHC positive control:	adenocarcinoma of human ovary tissue and human pancreas tissue; adenocarcinoma of human endometrium tissue and a denocarcinoma of human colon tissue
IHC Recommend dilution:	30-150
WB Predicted band size:	39 kDa
WB Positive control:	HepG2, Hela, SVT2, A549, COS7, PC12, MCF-7 cell lysates
WB Recommended dilution:	500-2000