

兔抗 CTDSP1 多克隆抗体

中文名称：兔抗 CTDSP1 多克隆抗体

英文名称：Anti-CTDSP1 rabbit polyclonal antibody

别名：NIF3; SCP1; NLIIF; NLI-IF

抗原：CTDSP1

储存：冷冻（-20℃）避光

宿主：Rabbit

反应种属：Human

相关类别：一抗

标记物：Unconjugate

克隆类型：Unconjugate

技术规格

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Background: | This gene encodes a member of the small C-terminal domain phosphatase (SCP) family of nuclear phosphatases. These proteins play a role in transcriptional regulation through specific dephosphorylation of phosphoserine 5 within tandem heptapeptide repeats of the C-terminal domain of RNA polymerase II. The encoded protein plays a role in neuronal gene silencing in non-neuronal cells, and may also inhibit osteoblast differentiation. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. |
| Applications: | WB, IHC |
| Name of antibody: | CTDSP1 |
| Immunogen: | Synthesized peptide derived from N-terminal of human CTD |

| | |
|---------------------------------|-------------------------------------------------------------------------------------|
| | SP1. |
| Full name: | CTD (carboxy-terminal domain, RNA polymerase II, polypeptide A) small phosphatase 1 |
| Synonyms : | NIF3; SCP1; NLIIF; NLI-IF |
| SwissProt: | Q9GZU7 |
| IHC positive control: | Human breast carcinoma tissue |
| IHC Recommend dilution: | 50-100 |
| WB Predicted band size: | 29 kDa |
| WB Positive control: | Huvec cells and Hela cells lysates |
| WB Recommended dilution: | 500-3000 |



