

兔抗 CYP2D6 多克隆抗体

中文名称：兔抗 CYP2D6 多克隆抗体

英文名称：Anti-CYP2D6 rabbit polyclonal antibody

别名：CPD6; CYP2D; CYP2DL1; CYP2D6; P450C2D; P450DB1; CYP2D7A; CYP2D7B; CYP2D7P; CYP2D8P; P450-DB1

相关类别：一抗

储存：冷冻 (-20℃)

宿主：Rabbit

抗原：CYP2D6

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and is known to metabolize as many as 20% of commonly prescribed drugs. Its substrates include debrisoquine, an adrenergic-blocking drug; sparteine and propafenone, both anti-arrhythmic drugs; and amitriptyline, an anti-depressant.
Applications:	ELISA, WB, IHC
Name of antibody:	CYP2D6

Immunogen:	Synthetic peptide of human CYP2D6
Full name:	cytochrome P450, family 2, subfamily D, polypeptide 6
Synonyms:	CPD6; CYP2D; CYP2DL1; CYP2D6; P450C2D; P450DB1; CYP2D7A; CYP2D7B; CYP2D7P2; CYP2D8P2; P450-DB1
SwissProt:	P10635
IHC positive control:	Human gastric cancer and human thyroid cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	56 kDa
WB Positive control:	K562 cells
WB Recommended dilution:	200-1000



