

DHX58 抗原（重组蛋白）

中文名称： DHX58 抗原（重组蛋白）

英文名称： DHX58 Antigen (Recombinant Protein)

别名： LGP2; RLR-3; D11LGP2; D11lgp2e

储存： 冷冻（-20℃）

相关类别： 抗原

概述：

Fusion protein corresponding to a region derived from 479-678 amino acids of human DHX58

技术规格：

Full name:	DExH-box helicase 58
Synonyms:	LGP2; RLR-3; D11LGP2; D11lgp2e
Swissprot:	Q96C10
Gene Accession:	BC014949
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	Acts as a regulator of DDX58/RIG-I and IFIH1/MDA5 mediated antiviral signaling. Cannot initiate antiviral signaling as it lacks the CARD domain required for activating MAVS/IPS1-dependent signaling events. Can have both negative and positive regulatory functions related to DDX58/RIG-I and IFIH1/MDA5 signaling and this role in regulating signaling may be complex and could probably depend on characteristics of the infecting virus or target cells, or both. Its inhibitory action on DDX58/RIG-I signaling may involve the following mechanisms: competition with DDX58/RIG-I for binding to the viral RNA, binding to DDX58/RIG-I and inhibiting its dimerization and interaction with MAVS/IPS1, competing with IKKε in its binding to MAVS/IPS1 thereby inhibiting activation

n of interferon regulatory factor 3 (IRF3). Its positive regulatory role may involve unwinding or stripping nucleoproteins of viral RNA thereby facilitating their recognition by DDX58/RIG-I and IFIH1/MDA5. Involved in the innate immune response to various RNA viruses and some DNA viruses such as poxviruses, and also to the bacterial pathogen *Listeria monocytogenes*. Can bind both ssRNA and dsRNA, with a higher affinity for dsRNA. Shows a preference to 5'-triphosphorylated RNA, although it can recognize RNA lacking a 5'-triphosphate.