

HLA-DPB1 抗原（重组蛋白）

中文名称：HLA-DPB1 抗原（重组蛋白）

英文名称：HLA-DPB1 Antigen (Recombinant Protein)

别名：major histocompatibility complex, class II, DP beta 1; DPB1; HLA-DP; HLA-DPB; HLA-DP1B

储存：冷冻（-20℃）

相关类别：抗原

概述：

Fusion protein corresponding to a region derived from 29-257 amino acids of human HLA-DPB1

技术规格：

| | |
|---------------------------|---|
| Full name: | major histocompatibility complex, class II, DP beta 1 |
| Synonyms: | DPB1; HLA-DP; HLA-DPB; HLA-DP1B |
| Swissprot: | P04440 |
| Gene Accession: | BC015000 |
| Purity: | >85%, as determined by Coomassie blue stained SDS-PAGE |
| Expression system: | Escherichia coli |
| Tags: | His tag C-Terminus, GST tag N-Terminus |
| Background: | HLA-DPB belongs to the HLA class II beta chain paralogues. This class II molecule is a heterodimer consisting of an alpha (DP A) and a beta chain (DPB), both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains. |

s, exon 4 encodes the transmembrane domain and exon 5 encodes the cytoplasmic tail. Within the DP molecule both the alpha chain and the beta chain contain the polymorphisms specifying the peptide binding specificities, resulting in up to 4 different molecules.