

兔抗 MID1 多克隆抗体

- 中文名称: 兔抗 MID1 多克隆抗体
- 英文名称: Anti-MID1 rabbit polyclonal antibody
- 别名: OS; FXY; OSX; OGS1; XPRF; BBBG1; GBBB1; MIDIN; RNF59; ZNFXY; TRIM18
- 相关类别: 一抗
- 储存: 冷冻(-20℃) 避光
- 抗原: MID1
- 宿 主: Rabbit
- 反应种属: Human Mouse Rat
- 标记物: Unconjugate
- 克隆类型: Unconjugate


技术规格

	The protein encoded by this gene is a member of the tripa rtite motif (TRIM) family, also known as the 'RING-B box-coi led coil' (RBCC) subgroup of RING finger proteins. The TRI
	M motif includes three zinc-binding domains, a RING, a B-b
	ox type 1 and a B-box type 2, and a coiled-coil region. Thi
Background:	s protein forms homodimers which associate with microtubu
	les in the cytoplasm. The protein is likely involved in the fo
	rmation of multiprotein structures acting as anchor points t
	o microtubules. Mutations in this gene have been associate
	d with the X-linked form of Opitz syndrome, which is chara
	cterized by midline abnormalities such as cleft lip, laryngeal





	cleft, heart defects, hypospadias, and agenesis of the corpus callosum.
Applications:	WB, IHC, IF
Name of antibody:	MID1
Immunogen:	Synthesized peptide derived from internal of human TRI18.
Full name:	midline 1
Synonyms :	OS; FXY; OSX; OGS1; XPRF; BBBG1; GBBB1; MIDIN; RNF59; Z NFXY; TRIM18
SwissProt:	015344
IHC positive control:	Human brain tissue
IHC Recommend dilution:	50-100
WB Predicted band size:	75 kDa
WB Positive control:	293 cells lysate
WB Recommended dilution:	500-3000
IF positive control:	Hela cells
IF Recommend dilution:	100-500





全国订货电话 4008-723-722

