

## 兔抗 MRC2 多克隆抗体

中文名称: 兔抗 MRC2 多克隆抗体

英文名称: Anti-MRC2 rabbit polyclonal antibody

别名: CD280; UPARAP; CLEC13E; ENDO180

抗原: MRC2

储存: 冷冻 (-20℃) 避光

宿主: Rabbit

相关类别: 一抗

反应种属: Human Mouse Rat

标记物: Unconjugate

克隆类型: Unconjugate

### 技术规格

|                          |  |
|--------------------------|--|
| <b>Background:</b>       | This gene encodes a member of the mannose receptor family of proteins that contain a fibronectin type II domain and multiple C-type lectin-like domains. The encoded protein plays a role in extracellular matrix remodeling by mediating the internalization and lysosomal degradation of collagen ligands. Expression of this gene may play a role in the tumorigenesis and metastasis of several malignancies including breast cancer, gliomas and metastatic bone disease. |
| <b>Applications:</b>     | WB, IHC, IF  |
| <b>Name of antibody:</b> | MRC2   |
| <b>Immunogen:</b>        | Synthesized peptide derived from internal of human MRC2.   |
| <b>Full name:</b>        | mannose receptor, C type 2   |
| <b>Synonyms :</b>        | CD280; UPARAP; CLEC13E; ENDO180  |
| <b>SwissProt:</b>        | Q9UBG0   |

|                                 |                    |
|---------------------------------|--------------------|
| <b>IHC positive control:</b>    | Human brain tissue |
| <b>IHC Recommend dilution:</b>  | 50-100             |
| <b>WB Predicted band size:</b>  | 167 kDa            |
| <b>WB Positive control:</b>     | 293 cells lysate   |
| <b>WB Recommended dilution:</b> | 500-3000           |
| <b>IF positive control:</b>     | HepG2 cells        |
| <b>IF Recommend dilution:</b>   | 100-500            |



