

兔抗 MUC1(CT)多克隆抗体

中文名称: 兔抗 MUC1(CT)多克隆抗体

英文名称: Anti-MUC1(CT) rabbit polyclonal antibody

别 名: EMA; MCD; PEM; PUM; KL-6; MAM6; MCKD; PEMT; CD227; H23AG; MCKD1; MUC-1; ADMCKD; ADMCKD1; CA 15-3; MUC-1/X; MUC1/ZD; MUC-1/SEC

- 抗原: MUC1(CT)
- 储存: 冷冻(-20℃)
- 宿 主: Rabbit
- 相关类别: 一抗
- 反应种属: Human, Mouse
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

ctions in cell-adhesion and the C-terminal beta subunit is involved in cell signaling. Overexpression, aberrant intrace	Background:	This gene encodes a membrane-bound protein that is a member of the mucin family. Mucins are O-glycosylated proteins that play an essential role in forming protective mucous barriers on epithelial surfaces. These proteins als o play a role in intracellular signaling. This protein is exp ressed on the apical surface of epithelial cells that line t he mucosal surfaces of many different tissues including I ung, breast stomach and pancreas. This protein is proteo lytically cleaved into alpha and beta subunits that form a heterodimeric complex. The N-terminal alpha subunit fun
Ilular localization, and changes in glycosylation of this pr		ctions in cell-adhesion and the C-terminal beta subunit is involved in cell signaling. Overexpression, aberrant intrace



全国订货电话 4008-723-722

	otein have been associated with carcinomas. This gene is known to contain a highly polymorphic variable number tandem repeats (VNTR) domain. Alternate splicing results in multiple transcript variants.
Applications:	ELISA, IHC
Name of antibody:	MUC1(CT)
Immunogen:	Synthetic peptide of human MUC1(CT)
Full name:	mucin 1, cell surface associated
Synonyms :	EMA; MCD; PEM; PUM; KL-6; MAM6; MCKD; PEMT; CD22 7; H23AG; MCKD1; MUC-1; ADMCKD; ADMCKD1; CA 15- 3; MUC-1/X; MUC1/ZD; MUC-1/SEC
SwissProt:	P15941
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human breast cancer and Human esophagus cancer
IHC Recommend dilution:	25-100

