

兔抗 FKBP3 多克隆抗体

| | |
|-------|--|
| 中文名称: | 兔抗 FKBP3 多克隆抗体 |
| 英文名称: | Anti-FKBP3 rabbit polyclonal antibody |
| 别名: | FKBP prolyl isomerase 3; FKBP-3; FKBP25; PPIase; FKBP-25 |
| 相关类别: | 一抗 |
| 储存: | 冷冻 (-20℃) |
| 宿主: | Rabbit |
| 抗原: | FKBP3 |
| 反应种属: | Human, Mouse |
| 标记物: | Unconjugate |
| 克隆类型: | rabbit polyclonal |

技术规格

| | |
|--------------------------|---|
| Background: | The protein encoded by this gene is a member of the immunophilin protein family, which play a role in immunoregulation and basic cellular processes involving protein folding and trafficking. This encoded protein is a cis-trans prolyl isomerase that binds the immunosuppressants FK506 and rapamycin, as well as histone deacetylases, the transcription factor YY1, casein kinase II, and nucleolin. It has a higher affinity for rapamycin than for FK506 and thus may be an important target molecule for immunosuppression by rapamycin. |
| Applications: | ELISA, WB, IHC |
| Name of antibody: | FKBP3 |
| Immunogen: | Fusion protein of human FKBP3 |

| | |
|------------------------------------|---|
| Full name: | FKBP prolyl isomerase 3 |
| Synonyms: | FKBP-3; FKBP25; PPIase; FKBP-25 |
| SwissProt: | Q00688 |
| ELISA Recommended dilution: | 5000-10000 |
| IHC positive control: | Human liver cancer |
| IHC Recommend dilution: | 200-300 |
| WB Predicted band size: | 25 kDa |
| WB Positive control: | Human bladder transitional cell carcinoma grade 2-3 tissue, Human bladder transitional cell carcinoma cancer next level 2-3 tissue lysate |
| WB Recommended dilution: | 1000-5000 |



