

## 兔抗 MAP3K5(Ab-966)多克隆抗体

中文名称： 兔抗 MAP3K5(Ab-966)多克隆抗体

英文名称： Anti-MAP3K5(Ab-966) rabbit polyclonal antibody

别名： ASK1; MEKK5; MAPKKK5

相关类别： 一抗

储存： 冷冻（-20℃）

抗原： MAP3K5(Ab-966)

宿主： Rabbit

反应种属： Human, Mouse, Rat

标记物： Unconjugate

克隆类型： rabbit polyclonal

### 技术规格

<b>IF Recommended dilution</b>	100-200
<b>IF Positive control:</b>	Hela cells
<b>WB Recommended dilution:</b>	500-1000
<b>WB Positive control:</b>	293 and Hela cell lysates
<b>WB Predicted band size:</b>	155 kDa
<b>IHC Recommend dilution:</b>	50-100
<b>IHC positive control:</b>	Human breast carcinoma
<b>SwissProt:</b>	Q99683
<b>Synonyms:</b>	ASK1; MEKK5; MAPKKK5
<b>Full name:</b>	mitogen-activated protein kinase kinase kinase 5
<b>Immunogen:</b>	Peptide sequence around aa.964-968 (S-I-S-L-P) derived from Human MAP3K5
<b>Name of antibody:</b>	MAP3K5(Ab-966)
<b>Applications:</b>	WB, IHC, IF
<b>Background:</b>	Mitogen-activated protein kinase (MAPK) signaling cascades include MAPK or extracellular signal-regulated kinase (ERK),

MAPK kinase (MKK or MEK), and MAPK kinase kinase (MAPKKK or MEKK). MAPKK kinase/MEKK phosphorylates and activates its downstream protein kinase, MAPK kinase/MEK, which in turn activates MAPK. The kinases of these signaling cascades are highly conserved, and homologs exist in yeast, Drosophila, and mammalian cells. MAPKKK5 contains 1,374 amino acids with all 11 kinase subdomains. Northern blot analysis shows that MAPKKK5 transcript is abundantly expressed in human heart and pancreas. The MAPKKK5 protein phosphorylates and activates MKK4 (aliases SERK1, MAPK K4) in vitro, and activates c-Jun N-terminal kinase (JNK)/stress-activated protein kinase (SAPK) during transient expression in COS and 293 cells; MAPKKK5 does not activate MAPK/ERK.



