

兔抗 MORF4L2 多克隆抗体

中文名称：兔抗 MORF4L2 多克隆抗体

英文名称：Anti-MORF4L2 rabbit polyclonal antibody

别名：mortality factor 4 like 2; MRGX; MORFL2

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：MORF4L2

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

The members of the mortality factor family include mortality factor 4 (MORF4), mortality factor 4 like 1 (MORF4L1), also known as MORF4-related gene 15 (MRG15), and mortality factor 4 like 2 (MORF4L2), also known as MORF-4 related gene X (MRGX). The human MORF4 gene maps to chromosome 4q34.1. MORF4 induces a senescent-like phenotype in complementation group B immortal cell lines. The genes encoding MORF4L1 and MORF4L2 map to chromosomes 15q25.1 and Xq22.2, respectively. MORF4, MORF4L1 and MORF4L2 each contain a C-terminal leucine zipper. An association between MORF4L1, Rb (retinoblastoma tumor suppressor) and PAM14 (protein associated with MORF4L1) suggests a role for MORF4L1 in transcription regulation. MORF4L1 also associates with the histone acetyl transferase MOF. In addition, MORF4, M

| | |
|------------------------------------|---|
| | ORF4L1 and MORF4L2 interact with mSin3A and TLE (transducin-like enhancer of split). The MORF/mSin3A/TLE association may repress transcription. In Purkinje cells, MORF4L1 localizes to the dendrites and the nuclei. |
| Applications: | ELISA, IHC |
| Name of antibody: | MORF4L2 |
| Immunogen: | Synthetic peptide of human MORF4L2 |
| Full name: | mortality factor 4 like 2 |
| Synonyms: | MRGX; MORFL2 |
| SwissProt: | Q15014 |
| ELISA Recommended dilution: | 5000-10000 |
| IHC positive control: | Human gastric cancer |
| IHC Recommend dilution: | 25-100 |



