

兔抗 PTK2 (phospho-Tyr576/Tyr577)多克隆抗体

中文名称： 兔抗 PTK2 (phospho-Tyr576/Tyr577)多克隆抗体

英文名称： Anti-PTK2 (phospho-Tyr576/Tyr577) rabbit polyclonal antibody

别 名： FAK; FADK; FAK1; FRNK; PPP1R71; p125FAK; pp125FAK

相关类别： 一抗

储 存： 冷冻 (-20°C) 避光

宿 主： Rabbit

抗 原： PTK2 (phospho-Tyr576/Tyr577)

反应种属： Human, Mouse, Rat

标 记 物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:	Non-receptor protein-tyrosine kinase implicated in signaling pathways involved in cell motility, proliferation and apoptosis. Activated by tyrosine-phosphorylation in response to either integrin clustering induced by cell adhesion or antibody cross-linking, or via G-protein coupled receptor (GPCR) occupancy by ligands such as bombesin or lysophosphatidic acid, or via LDL receptor occupancy. Plays a potential role in oncogenic transformations resulting in increased kinase activity.
Applications:	WB, IF
Name of antibody:	PTK2 (phospho-Tyr576/Tyr577)

Immunogen:	Synthetic peptide of human PTK2 (phospho-Tyr576/Tyr577)
Full name:	protein tyrosine kinase 2 (phospho-Tyr576/Tyr577)
Synonyms :	FAK; FAKD; FAK1; FRNK; PPP1R71; p125FAK; pp125FAK
SwissProt:	Q05397
WB Predicted band size:	125 kDa
WB Positive control:	HT29 cells treated with PMA
WB Recommended dilution:	500-1000
IF Positive control:	HeLa cells
IF Recommended dilution	100-200



