

兔抗 MAF 多克隆抗体

中文名称: 兔抗 MAF 多克隆抗体

英文名称: Anti-MAF rabbit polyclonal antibody

别 名: CCA4; c-MAF

相关类别: 一抗

储 存: 冷冻 (-20℃)

宿 主: Rabbit

抗 原: MAF

反应种属: Human, Mouse, Rat

标 记 物: Unconjugate

克隆类型: rabbit polyclonal


技术规格

Background:	The protein encoded by this gene is a DNA-binding, leuci ne zipper-containing transcription factor that acts as a ho modimer or as a heterodimer. Depending on the binding site and binding partner, the encoded protein can be a transcriptional activator or repressor. This protein plays a role in the regulation of several cellular processes, including embryonic lens fiber cell development, increased T-cell sus ceptibility to apoptosis, and chondrocyte terminal different iation. Defects in this gene are a cause of juvenile-onset pulverulent cataract as well as congenital cerulean cataract 4 (CCA4). Two transcript variants encoding different isoforms have been found for this gene.
Applications:	ELISA, WB, IHC
Name of antibody:	MAF




全国订货电话 4008-723-722

Immunogen:	Synthetic peptide of human MAF
Full name:	MAF bZIP transcription factor
Synonyms:	CCA4; AYGRP; c-MAF; CTRCT21
SwissProt:	O75444
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human ovarian cancer
IHC Recommend dilution:	50-100
WB Predicted band size:	38 kDa
WB Positive control:	HepG2 cell and Mouse brain tissue lysates
WB Recommended dilution:	500-2000









全国订货电话 4008-723-722