

兔抗 SMC1A 多克隆抗体

- 中文名称：兔抗 SMC1A 多克隆抗体
- 英文名称：Anti-SMC1A rabbit polyclonal antibody
- 别名：SMC1; SMCB; CDLS2; SB1.8; SMC1L1; DXS423E; SMC1alpha
- 抗原：SMC1A
- 储存：冷冻 (-20℃)
- 宿主：Rabbit
- 相关类别：一抗
- 反应种属：Human, Mouse, Rat
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

技术规格

Background:	Proper cohesion of sister chromatids is a prerequisite for the correct segregation of chromosomes during cell division. The cohesin multiprotein complex is required for sister chromatid cohesion. This complex is composed partly of two structural maintenance of chromosomes (SMC) proteins, SMC3 and either SMC1B or the protein encoded by this gene. Most of the cohesin complexes dissociate from the chromosomes before mitosis, although those complexes at the kinetochore remain. Therefore, the encoded protein is thought to be an important part of functional kinetochores. In addition, this protein interacts with BRCA1 an
--------------------	---

	<p>d is phosphorylated by ATM, indicating a potential role for this protein in DNA repair. This gene, which belongs to the SMC gene family, is located in an area of the X-chromosome that escapes X inactivation. Mutations in this gene result in Cornelia de Lange syndrome. Alternative splicing results in multiple transcript variants encoding different isoforms.</p>
Applications:	ELISA, IHC
Name of antibody:	SMC1A
Immunogen:	Fusion protein of human SMC1A
Full name:	structural maintenance of chromosomes 1A
Synonyms:	SMC1; SMCB; CDLS2; SB1.8; SMC1L1; DXS423E; SMC1alpha
SwissProt:	Q14683
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human breast cancer and human esophagus cancer
IHC Recommend dilution:	30-150



