

兔抗 STARD13 多克隆抗体

- 中文名称: 兔抗 STARD13 多克隆抗体
- 英文名称: Anti-STARD13 rabbit polyclonal antibody
- 别名: DLC2; GT650; ARHGAP37; LINC00464
- 抗 原: STARD13
- 储存: 冷冻 (-20℃) 避光
- 宿 主: Rabbit
- 反应种属: Human
- 相关类别: 一抗
- 标记物: Unconjugate
- 克隆类型: Unconjugate

技术规格

Background:	This gene encodes a protein which contains an N-terminal sterile alpha motif (SAM) for protein-protein interactions, f ollowed by an ATP/GTP-binding motif, a GTPase-activating protein (GAP) domain, and a C-terminal STAR-related lipid transfer (START) domain. It may be involved in regulation of cytoskeletal reorganization, cell proliferation, and cell m otility, and acts as a tumor suppressor in hepatoma cells. The gene is located in a region of chromosome 13 that is associated with loss of heterozygosity in hepatocellular car cinomas. Alternatively spliced transcript variants encoding
	cinomas. Alternatively spliced transcript variants encoding different isoforms have been described for this gene.



Applications:	WB, IF
Name of antibody:	STARD13
Immunogen:	Synthesized peptide derived from N-terminal of human ST A13.
Full name:	StAR-related lipid transfer (START) domain containing 13
Synonyms :	DLC2; GT650; ARHGAP37; LINC00464
SwissProt:	Q9Y3M8
WB Predicted band size:	125 kDa
WB Positive control:	K562 cells lysate
WB Recommended dilution:	500-3000
IF positive control:	A549 cells
IF Recommend dilution:	100-500





