

兔抗 TACC2 多克隆抗体

- 中文名称：兔抗 TACC2 多克隆抗体
- 英文名称：Anti-TACC2 rabbit polyclonal antibody
- 别名：AZU-1; ECTACC
- 相关类别：一抗
- 储存：冷冻（-20℃）
- 宿主：Rabbit
- 抗原：TACC2
- 反应种属：Human, Mouse
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

技术规格

Background:

Transforming acidic coiled-coil proteins are a conserved family of centrosome- and microtubule-interacting proteins that are implicated in cancer. This gene encodes a protein that concentrates at centrosomes throughout the cell cycle. This gene lies within a chromosomal region associated with tumorigenesis. Expression of this gene is induced by erythropoietin and is thought to affect the progression of breast tumors. Several transcript variants encoding different isoforms have been found for this gene.

Applications:	ELISA, IHC
Name of antibody:	TACC2
Immunogen:	Synthetic peptide of human TACC2
Full name:	transforming, acidic coiled-coil containing protein 2
Synonyms :	AZU-1; ECTACC
SwissProt:	O95359
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human liver cancer
IHC Recommend dilution:	25-100

