

兔抗 TNFRSF10C 多克隆抗体

- 中文名称：兔抗 TNFRSF10C 多克隆抗体
- 英文名称：Anti-TNFRSF10C rabbit polyclonal antibody
- 别名：LIT; DCR1; TRID; CD263; TRAILR3; TRAIL-R3; DCR1-TNFR
- 抗原：TNFRSF10C
- 储存：冷冻（-20℃）
- 宿主：Rabbit
- 相关类别：一抗
- 反应种属：Human
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

技术规格

Background:

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor contains an extracellular TRAIL-binding domain and a transmembrane domain, but no cytoplasmic death domain. This receptor is not capable of inducing apoptosis, and is thought to function as an antagonistic receptor that protects cells from TRAIL-induced apoptosis. This gene was found to be a p53-regulated DNA damage-inducible gene. The expression of this gene was detected in many normal tissues but not in most cancer cell lines, which m

	ay explain the specific sensitivity of cancer cells to the apoptosis-inducing activity of TRAIL.
Applications:	ELISA, IHC
Name of antibody:	TNFRSF10C
Immunogen:	Fusion protein of human TNFRSF10C
Full name:	tumor necrosis factor receptor superfamily member 10c
Synonyms:	LIT; DCR1; TRID; CD263; TRAILR3; TRAIL-R3; DCR1-TNFR
SwissProt:	O14798
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human liver cancer and human lung cancer
IHC Recommend dilution:	25-100



