

兔抗 TRAF3 多克隆抗体

- 中文名称: 兔抗 TRAF3 多克隆抗体
- 英文名称: Anti-TRAF3 rabbit polyclonal antibody
- 别名: CAP1, LAP1, CAP-1, CRAF1, IIAE5, CD40bp
- 相关类别: 一抗
- 储存: 冷冻(-20℃)
- 宿 主: Rabbit
- 抗 原: TRAF3
- 反应种属: Human, Mouse
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

NF-kappaß activation and cell death initiated by Libeta	Background:	The protein encoded by this gene is a member of the TNF receptor associated factor (TRAF) protein family. T RAF proteins associate with, and mediate the signal tra nsduction from, members of the TNF receptor (TNFR) s uperfamily. This protein participates in the signal transd uction of CD40, a TNFR family member important for t he activation of the immune response. This protein is f ound to be a critical component of the lymphotoxin-be ta receptor (LTbetaR) signaling complex, which induces
ligation. Epstein-Barr virus encoded latent infection me		NF-kappaB activation and cell death initiated by LTbeta



	mbrane protein-1 (LMP1) can interact with this and sev eral other members of the TRAF family, which may be essential for the oncogenic effects of LMP1. Several alt ernatively spliced transcript variants encoding three dist inct isoforms have been reported.
Applications:	ELISA, WB, IHC
Name of antibody:	TRAF3
Immunogen:	Synthetic peptide of human TRAF3
Full name:	TNF receptor-associated factor 3
Synonyms :	CAP1, LAP1, CAP-1, CRAF1, IIAE5, CD40bp
SwissProt:	Q13114
ELISA Recommended dilution:	1000-5000
IHC positive control:	Human tonsil
IHC Recommend dilution:	25-100
WB Predicted band size:	64 kDa
WB Positive control:	Mouse kidney tissue
WB Recommended dilution:	200-1000





