

兔抗 BCL7A 多克隆抗体

中文名称: 兔抗 BCL7A 多克隆抗体

英文名称: Anti-BCL7A rabbit polyclonal antibody

别名: BCL7

相关类别: 一抗

储存: 冷冻 (-20℃) 避光

宿主: Rabbit

抗原: BCL7A

反应种属: Human

标记物: Unconjugate

克隆类型: Unconjugate

技术规格

Background:	This gene is directly involved, with Myc and IgH, in a t hree-way gene translocation in a Burkitt lymphoma cell line. As a result of the gene translocation, the N-termin al region of the gene product is disrupted, which is th ought to be related to the pathogenesis of a subset of high-grade B cell non-Hodgkin lymphoma. The N-termin al segment involved in the translocation includes the r egion that shares a strong sequence similarity with tho se of BCL7B and BCL7C. Two transcript variants encodi ng different isoforms have been found for this gene.
Applications:	WB
Name of antibody:	BCL7A
Immunogen:	Synthesized peptide derived from internal of human BC L7A.

Full name:	B-cell CLL/lymphoma 7A
Synonyms :	BCL7
SwissProt:	Q4VC05
WB Predicted band size:	23 kDa
WB Positive control:	Huvec cells lysate
WB Recommended dilution:	500-3000

