

兔抗 CD207 多克隆抗体

中文名称: 兔抗 CD207 多克隆抗体

英文名称: Anti-CD207 rabbit polyclonal antibody

别 名: CLEC4K

抗 原: CD207

储 存: 冷冻(-20℃)

宿 主: Rabbit

相关类别: 一抗

反应种属: Human, Mouse

标记物: Unconjugate

克隆类型: rabbit polyclonal


技术规格

Background:	The protein encoded by this gene is expressed only in Langerhans cells which are immature dendritic cell s of the epidermis and mucosa. It is localized in the Birbeck granules, organelles present in the cytoplasm of Langerhans cells and consisting of superimposed and zippered membranes. It is a C-type lectin with mannose binding specificity, and it has been propos ed that mannose binding by this protein leads to int ernalization of antigen into Birbeck granules and pro viding access to a nonclassical antigen-processing pa thway. Mutations in this gene result in Birbeck granules deficiency or loss of sugar binding activity.
Applications:	ELISA, WB, IHC
Name of antibody:	CD207



全国订货电话 4008-723-722

Immunogen:	Fusion protein of human CD207
Full name:	CD207 molecule
Synonyms:	CLEC4K
SwissProt:	Q9UJ71
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer
IHC Recommend dilution:	250-500
WB Predicted band size:	37 kDa
WB Positive control:	Mouse skin tissue lysate
WB Recommended dilution:	500-2000





全国订货电话 4008-723-722

